-continued

```
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<400> SEQUENCE: 100
aaataccctt acatccataa
                                                                        20
<210> SEQ ID NO 101
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE
<223> OTHER INFORMATION: Synthetic oligonucleotide
<400> SEOUENCE: 101
accettacat ccataat
                                                                        17
<210> SEQ ID NO 102
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<400> SEOUENCE: 102
                                                                        18
ccttacatcc ataatcat
<210> SEQ ID NO 103
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic oligonucleotide
<400> SEQUENCE: 103
cttacatcca taatcattt
                                                                        19
```

- 1. An antisense oligonucleotide, 10-30 nucleotides in length, wherein said antisense oligonucleotide comprises a contiguous nucleotide sequence 10-30 nucleotides in length, wherein the contiguous nucleotide sequence is at least 90% complementary, such as fully complementary to SEQ ID NO 1, wherein the antisense oligonucleotide is capable of inhibiting the expression of human TIA1 in a cell which is expressing human TIA1; or a pharmaceutically acceptable salt thereof.
- 2. The antisense oligonucleotide according to claim 1, wherein the contiguous nucleotide sequence is at least 90% complementary, such as fully complementary to a sequence selected from the group consisting of SEQ ID NO 4-53.
- **3**. The antisense oligonucleotide according to any one of claims **1** to **3**, wherein the contiguous nucleotide sequence is fully complementary to a region of SEQ ID NO 1, selected from the group consisting of the regions in list A.
- 4. The antisense oligonucleotide according to any one of claims 1 to 3, wherein the contiguous nucleotide sequence is fully complementary to a region of SEQ ID NO 1, selected from the group consisting of the regions in list B.
- **5**. The antisense oligonucleotide according to any one of claims **1-4**, wherein the antisense oligonucleotide is a gapmer oligonucleotide comprising a contiguous nucleotide sequence of formula 5'-F-G-F'-3', where region F and F' independently comprise 1-8 sugar modified nucleosides, and

- G is a region between 5 and 16 nucleosides which are capable of recruiting RNaseH.
- **6.** The antisense oligonucleotide according to claim **5**, wherein the sugar modified nucleosides of region F and F' are independently selected from the group consisting of 2'-O-alkyl-RNA, 2'-O-methyl-RNA, 2'-alkoxy-RNA, 2'-O-methoxyethyl-RNA, 2'-amino-DNA, 2'-fluoro-DNA, arabino nucleic acid (ANA), 2'-fluoro-ANA and LNA nucleosides.
- 7. The antisense oligonucleotide according to claim 5 or 6, wherein region G comprises 5-16 contiguous DNA nucleosides.
- **8**. The antisense oligonucleotide according to any one of claims 1-7, wherein the antisense oligonucleotide is a LNA gapmer oligonucleotide.
- 9. The antisense oligonucleotide according to any one of claims 5-8, wherein the LNA nucleosides are beta-D-oxy LNA nucleosides.
- 10. The antisense oligonucleotide according to any one of claims 1-9, wherein the internucleoside linkages between the contiguous nucleotide sequence are phosphorothioate internucleoside linkages.
- 11. The antisense oligonucleotide according to any one of claims 1-10, wherein the oligonucleotide comprises a contiguous nucleotide sequence selected from the group consisting of: 54-103.